



Consent form for COVID-19 vaccination

Before you fill out this form, make sure you read the information sheet on the vaccine you will be getting: Vaxzevria (AstraZeneca), Comirnaty (Pfizer) or Spikevax (Moderna).

Last updated: 13 December 2021

About COVID-19 vaccination

People who have a COVID-19 vaccination have a much lower chance of getting sick from COVID-19.

There are three brands of vaccine in use in Australia. All are effective and safe.

You can have:

- AstraZeneca if you are 18 years or over
- Moderna if you are 12 years or over
- Pfizer if you are 5 years or over*

Pfizer or Moderna are preferred over AstraZeneca for adults under 60 years of age.

Most people require two doses initially. This is called the primary course.

People with severe immunocompromise may require a third primary dose to bring their immune response up to optimal levels

People aged 18 years or over can have a booster dose of Pfizer, Moderna (half dose), or AstraZeneca five months or more after their primary course, to prolong their protection against COVID-19.

For more information visit the Department of Health COVID-19 vaccine website: www.health.gov.au/covid-19-vaccines

Medical experts have studied COVID-19 vaccines to make sure they are safe. Most side effects are mild. They may start on the day of vaccination and last for one or two days. As with any vaccine or medicine, there may be rare or unknown side effects

Name:						
Medicare number:						

^{*}There is a separate consent form available for children aged 5 to 11 years.

A very rare side effect after AstraZeneca is called thrombosis with thrombocytopenia syndrome, or TTS. This means blood clotting (thrombosis) with low blood platelet levels (thrombocytopenia). TTS does not happen after Pfizer or Moderna.

Myocarditis and pericarditis (heart inflammation) have been reported following Pfizer and Moderna. Most cases have been mild and people have recovered quickly.

Tell your health care provider if you have any side effects after vaccination that you are worried about.

You may be contacted by SMS or email in the week after you have the vaccine to see how you are feeling.

Some people may get COVID-19 after vaccination. You must still follow all public health advice in your state or territory to stop the spread of COVID-19, including:

- keep your distance stay at least 1.5 metres away from other people
- wash your hands often with soap and water, or use hand sanitiser
- wear a mask
- stay home if you are unwell with cold or flu-like symptoms and arrange to get a COVID-19 test.

By law, the person giving your vaccination must record it on the Australian Immunisation Register. You can view your vaccination record online through your:

- Medicare account
- MyGov account
- My Health Record account (you can register for this with a Medicare number or an Individual Healthcare Identifier).

How your information is used

For information on how your personal details are collected, stored and used, visit www.health.gov.au/using-our-websites/privacy/privacy-notice-for-covid-19-vaccinations.

If you are getting your vaccination in a pharmacy, the pharmacy must share some of your personal information with the Pharmacy Programs Administrator. This is so the pharmacy can claim payment from the Australian Government. More information about why this is needed and what information is shared is provided at the link above.

On the day you have your vaccine

Before you get vaccinated, tell the person giving you the vaccination if you:

- have had an allergic reaction, particularly a severe allergic reaction (anaphylaxis), to:
 - o a previous dose of a COVID-19 vaccine
 - o an ingredient of a COVID-19 vaccine
 - o other vaccines or medications
- are immunocompromised. This means that you have a weakened immune system that
 makes it harder for you to fight diseases. You can still have a COVID-19 vaccine, but talk to
 your doctor about when is the best time to get your vaccine. This will depend on your
 condition and your treatment.

Name:						
Medicare number:						

Consent Checklist

get-vaccinated/pregnant-women

Yes	No	
		Have you had an allergic reaction to a previous dose of a COVID-19 vaccine?
		Have you had anaphylaxis to another vaccine or medication?
		Have you had a serious adverse event, that following expert review by an experienced immunisation provider or medical specialist was attributed to a previous dose of a COVID-19 vaccine (and did not have another cause identified)?
		Have you ever had mastocytosis (a mast cell disorder) which has caused recurrent anaphylaxis?
		Have you had COVID-19 before?
		Do you have a bleeding disorder?
		Do you take any medicine to thin your blood (an anticoagulant therapy)?
		Do you have a weakened immune system (immunocompromised)?
		Are you pregnant? *
		Have you been sick with a cough, sore throat, fever or are feeling sick in another way?
		Have you had a COVID-19 vaccination before?
		Have you received any other vaccination in the last 7 days?
Releva	ant on	ly for those receiving AstraZeneca:
		Have you ever been diagnosed with capillary leak syndrome?
		Have had thrombosis (clotting) together with thrombocytopenia (low platelets) within 42 days after having a previous dose of AstraZeneca?
		Have you ever had cerebral venous sinus thrombosis? *
		Have you ever had heparin-induced thrombocytopenia? *
		Have you ever had blood clots in the abdominal veins (splanchnic veins)? *
		Have you ever had antiphospholipid syndrome associated with blood clots? *
		Are you under 60 years of age? *
		lerna are the preferred vaccines for people in these groups. If these vaccines are not aZeneca can be considered if the benefits of vaccination outweigh the risks.
For more		mation, see https://www.health.gov.au/initiatives-and-programs/covid-19-vaccines/advice-s

Name:

Medicare number:

If you are pregnant, see https://www.health.gov.au/initiatives-and-programs/covid-19-vaccines/who-can-decomposition

Relevant onl	y for those receiving	Pfize	r or N	1oder	na:								
	Have you been diagnosed with myocarditis and/or pericarditis after a previous dose of Pfizer or Moderna?												
	Have you had myo	Have you had myocarditis or pericarditis within the past three months?											
	Do you currently ha	Do you currently have acute rheumatic fever or acute rheumatic heart disease?											
☐ ☐ Do you have severe heart failure?													
If you answered Yes to any of the above questions, you may still be able to receive Pfizer or Moderna, however you should talk to your GP, immunisation specialist or cardiologist first to discuss the best timing of vaccination and whether any additional precautions are needed. Last updated: 13 December 2021													
Patient information													
Name:													
Medicare no	umber:												
Individual H	lealth Identifier (IHI) e:										I		
Date of birth	า:												
Address:													
Phone cont	act number:												
Email addre	ess:												
Gender:													
Language s	poken at home:												
Country of b	oirth:												
Are you Abo	riginal and/or Torres	Strait	Islan	der?									
Are you Aboriginal and/or Torres Strait Islander? Yes, Aboriginal only Yes, Torres Strait Islander only Yes Aboriginal and Torres Strait Islander No Prefer not to answer													
Next of kin	(in case of emergeno	;y):											
Name:													
Phone cont	act number:												
			· <u></u>				· <u></u>						_

Name:						
Medicare number:						

Con	Consent to receive COVID-19 vaccine									
	I confirm I have revaccination.	eceived a	and understood information provided to me on COVID-19							
	I confirm that I have none of the above conditions apply to me, or I have discussed these conditions and any other special circumstances with my regular health care provider and/or vaccination provider.									
☐ I agree to receive a course of COVID-19 vaccine / I agree to receive a booster of COVID-19 vaccine										
Pati	ent's name:									
Pati	ent's signature:									
Date	э:									
	-	•	guardian or substitute decision-maker, and agree to the patient named above.							
	ent/guardian/subst sion-maker's nam									
	ent/guardian/subst sion maker's signa									
Date	e :									

Name:						
Medicare number:						

For provider use:

Dose 1:

2000	
Date vaccine administered:	
Time received:	
COVID-19 vaccine brand administered:	
Batch no:	
Serial no:	
Site of vaccine injection:	
Name of vaccination service provider:	
Dose 2:	
Date vaccine administered:	
Time received:	
COVID-19 vaccine brand administered:	
Batch no:	
Serial no:	
Site of vaccine injection:	
Name of vaccination service provider:	
Dose 3 or booster*:	
Date vaccine administered:	
Time received:	
COVID-19 vaccine brand administered:	
Batch no:	
Serial no:	
Site of vaccine injection:	
Name of vaccination service provider:	
	primary dose of COVID-19 vaccine in individuals I recommendations on the use of a booster dose ations on the use of Spikevax Moderna as a

booster dose.

Last updated: 13 December 2021

Name:						
Medicare number:						